


# X-ray diffraction


---

## Intensity of the scattered waves




$$F_1 \frac{\cos(k|\vec{r} - \vec{r}_1| - \omega t + kx_1)}{\sqrt{|\vec{r} - \vec{r}_1|}} + F_2 \frac{\cos(k|\vec{r} - \vec{r}_2| - \omega t + kx_2)}{\sqrt{|\vec{r} - \vec{r}_2|}}.$$

## Intensity of the scattered waves



$$\left( \frac{F_1}{\sqrt{|\vec{r} - \vec{r}_1|}} e^{i(k|\vec{r} - \vec{r}_1| + kx_1)} + \frac{F_2}{\sqrt{|\vec{r} - \vec{r}_2|}} e^{i(k|\vec{r} - \vec{r}_2| + kx_2)} \right) e^{-i\omega t}$$

# Interference



$$a = \frac{\vec{r} \cdot \vec{k}}{|\vec{k}|} \quad b = \frac{\vec{r} \cdot \vec{k}'}{|\vec{k}|}$$

elastic scattering


$$|k| = |k'|$$

path difference: 
$$a - b = \frac{-\vec{r} \cdot (\vec{k}' - \vec{k})}{|\vec{k}|}$$

phase shift: 
$$\varphi = 2\pi \frac{a - b}{\lambda} = 2\pi \frac{-\vec{r} \cdot (\vec{k}' - \vec{k})}{|\vec{k}| \lambda} = -\vec{r} \cdot (\vec{k}' - \vec{k}) = -\Delta\vec{k} \cdot \vec{r}$$

Amplitude: 
$$F = F_0 + F_0 e^{-i\Delta\vec{k} \cdot \vec{r}}$$


# Interference



Amplitude: 
$$F_{tot} = \sum_i F_i e^{-i\Delta\vec{k} \cdot \vec{r}_i}$$

# Diffraction


Scattering is  
proportional to electron  
density  $n(r)$



The scattering amplitude is proportional to the Fourier transform of the electron density.

# Scattering amplitude

---



$$F \propto \int n(\vec{r}) \exp(-i\Delta\vec{k} \cdot \vec{r}) dV$$

expand  $n(r)$  in a Fourier series

$$n(\vec{r}) = \sum_{\vec{G}} n_{\vec{G}} \exp(i\vec{G} \cdot \vec{r})$$

$$F \propto \sum_{\vec{G}} \int n_{\vec{G}} \exp(i(\vec{G} - \Delta\vec{k}) \cdot \vec{r}) dV$$

if  $\vec{G} = \Delta\vec{k}$ , all components add coherently


diffraction condition:  $\boxed{\vec{G} = \Delta\vec{k}}$

The intensity of the peak at  $G$  is  $|n_G|^2$

# Nobel Prize 1914

---

first diffraction experiment of Max von Laue 1912  
ZnS single crystal, exposure time 30'  
the 5<sup>th</sup> diffraction pattern



M. von Laue (1879-1960)


$$\vec{G} = \Delta \vec{k}$$



# Diffraction condition (Laue condition)

---


$$\vec{k}' - \vec{k} = \Delta\vec{k} = \vec{G}$$



$$|\vec{k}| = |\vec{k}'| \text{ for elastic scattering}$$

# Single crystal diffraction

---




$$\vec{G} = \Delta\vec{k} = \vec{q} \quad \leftarrow \text{Scattering wave vector}$$

Every time a diffraction peak is observed, record  $G$ . When many  $G$  vectors are known, determine the reciprocal lattice.

The sample and the detector must be turned to find all of the diffraction peaks.

Gx	Gy	Gz	$ n_G ^2$
2.4E10	2.4E10	0	10341
2.4E10	0	2.4E10	9989

$$n(\vec{r}) = \sum_{\vec{G}} n_{\vec{G}} \exp(i\vec{G} \cdot \vec{r})$$



$\theta$  sets the length of the scattering vector

# Determining real space primitive lattice vectors

---

$$\vec{a}_1 = 2\pi \frac{\vec{b}_2 \times \vec{b}_3}{\vec{b}_1 \cdot (\vec{b}_2 \times \vec{b}_3)}$$

$$\vec{a}_2 = 2\pi \frac{\vec{b}_3 \times \vec{b}_1}{\vec{b}_1 \cdot (\vec{b}_2 \times \vec{b}_3)}$$

$$\vec{a}_3 = 2\pi \frac{\vec{b}_1 \times \vec{b}_2}{\vec{b}_1 \cdot (\vec{b}_2 \times \vec{b}_3)}$$


$$\vec{a}_i \cdot \vec{b}_j = 2\pi \delta_{ij}$$

$\vec{b}_1, \vec{b}_2, \vec{b}_3$  determined  
from diffraction experiment

Volume of the primitive unit cell  $\vec{a}_1 \cdot (\vec{a}_2 \times \vec{a}_3)$

$G$  vectors specify the Bravais lattice.


# Ewald sphere $\Delta\vec{k} = \vec{G}$



Draw a vector representing the incoming radiation so that it ends at the origin. As the crystal is rotated around the origin, the condition for diffraction will be satisfied every time a reciprocal lattice point is on the sphere.

# Diffraction condition

---



$$\vec{k}' - \vec{k} = \vec{G}$$

For every  $G$  there is a  $-G$  so the diffraction condition can also be written as


$$\vec{k} - \vec{k}' = \vec{G}$$

a wave will be diffracted if the wave vector ends on one of the planes

# Brillouin zones




Leon Brillouin




1st Brillouin zone consists of the  $k$ -states around the origin that can be reached without crossing a plane.

# 1st Brillouin zones


---



sc



bcc










fcc

1st Brillouin is the Wigner-Seitz cell in reciprocal space.




## Brillouin zones

<p><b>Triclinic</b></p> <p><math>a \neq b \neq c</math>  <math>\alpha \neq \beta \neq \gamma</math></p>	 <p>Triclinic</p>	
<p><b>Monoclinic</b></p> <p><math>a \neq b \neq c</math>  <math>\alpha \neq 90^\circ</math>  <math>\beta = \gamma = 90^\circ</math></p>	 <p>Monoclinic simple</p>	 <p>Monoclinic Base centered</p>
<p><b>Orthorhombic</b></p> <p><math>a \neq b \neq c</math>  <math>\alpha = \beta = \gamma = 90^\circ</math></p>	 <p>Orthorhombic simple</p>	 <p>Base centered</p>
		

# Electron density of an atom

---



Most of the electrons are concentrated around the nucleus. The integral over the electron density is proportional to the number of electrons.

$$n_j(\vec{r}) \propto \exp\left(-\frac{(\vec{r} - \vec{r}_j)^2}{r_0^2}\right)$$

Approximately a Gaussian centered at  $r_j$

# Electron density

---

Write the electron density as a Fourier series

$$n(\vec{r}) = \sum_{\vec{G}} n_{\vec{G}} e^{i\vec{G} \cdot \vec{r}} = \sum_{\vec{T}} \sum_j n_j (\vec{r} - \vec{r}_j + \vec{T}),$$

position of atom  $j$  of the basis

Translation vector

Multiply by  $e^{-i\vec{G}' \cdot \vec{r}}$  and integrate over a unit cell.

$$\sum_{\vec{G}} \int_{\text{u.c.}} n_{\vec{G}} e^{i\vec{G} \cdot \vec{r}} e^{-i\vec{G}' \cdot \vec{r}} d\vec{r} = \sum_j \int_{\text{u.c.}} n_j (\vec{r} - \vec{r}_j) e^{-i\vec{G}' \cdot \vec{r}} d\vec{r}.$$

1


# Electron density

---

$$n_{\vec{G}} V = \sum_j \int n_j(\vec{r} - \vec{r}_j) e^{-i\vec{G} \cdot \vec{r}} d\vec{r}.$$

Make a substitution  $\vec{r}' = \vec{r} - \vec{r}_j$ .

$$n_{\vec{G}} = \frac{1}{V} \sum_j e^{-i\vec{G} \cdot \vec{r}_j} \underbrace{\int n_j(\vec{r}') e^{-i\vec{G} \cdot \vec{r}'} d\vec{r}'}.$$



Fourier transform of the electron density of atom  $j$   
= atomic form factor  $f_j(G)$

# Atomic form factor

---

$$f_j(\vec{G}) = \int n_j(\vec{r}) e^{-i\vec{G} \cdot \vec{r}} d\vec{r},$$

The atomic form factors can be looked up in a table.

The structure factors are given in terms of the atomic form factors.

$$n_{\vec{G}} = \frac{1}{V} \sum_j f_j(\vec{G}) e^{-i\vec{G} \cdot \vec{r}_j}$$

sum over the basis

position of atom  $j$  of the basis

## Section 4.3.2. Parameterizations of electron atomic scattering factors

J. M. Cowley,<sup>b†</sup> L. M. Peng,<sup>i</sup> G. Ren,<sup>j</sup> S. L. Dudarev<sup>c</sup> and M. J. Whelan<sup>c</sup>

**Table 4.3.2.2** | [pdf](#) |

Elastic atomic scattering factors of electrons for neutral atoms and  $s$  up to  $2.0 \text{ \AA}^{-1}$

$$f(s) = \sum_i a_i \exp(-b_i s^2)$$

Element	$Z$	$a_1$	$a_2$	$a_3$	$a_4$	$a_5$	$b_1$	$b_2$	$b_3$	$b_4$	$b_5$
H	1	0.0349	0.1201	0.1970	0.0573	0.1195	0.5347	3.5867	12.3471	18.9525	38.6269
He	2	0.0317	0.0838	0.1526	0.1334	0.0164	0.2507	1.4751	4.4938	12.6646	31.1653
Li	3	0.0750	0.2249	0.5548	1.4954	0.9354	0.3864	2.9383	15.3829	53.5545	138.7337
Be	4	0.0780	0.2210	0.6740	1.3867	0.6925	0.3131	2.2381	10.1517	30.9061	78.3273
B	5	0.0909	0.2551	0.7738	1.2136	0.4606	0.2995	2.1155	8.3816	24.1292	63.1314
C	6	0.0893	0.2563	0.7570	1.0487	0.3575	0.2465	1.7100	6.4094	18.6113	50.2523
N	7	0.1022	0.3219	0.7982	0.8197	0.1715	0.2451	1.7481	6.1925	17.3894	48.1431
O	8	0.0974	0.2921	0.6910	0.6990	0.2039	0.2067	1.3815	4.6943	12.7105	32.4726
F	9	0.1083	0.3175	0.6487	0.5846	0.1421	0.2057	1.3439	4.2788	11.3932	28.7881
Ne	10	0.1269	0.3535	0.5582	0.4674	0.1460	0.2200	1.3779	4.0203	9.4934	23.1278
Na	11	0.2142	0.6853	0.7692	1.6589	1.4482	0.3334	2.3446	10.0830	48.3037	138.2700
Mg	12	0.2314	0.6866	0.9677	2.1882	1.1339	0.3278	2.2720	10.9241	39.2898	101.9748
Al	13	0.2390	0.6573	1.2011	2.5586	1.2312	0.3138	2.1063	10.4163	34.4552	98.5344



## 513.001 Molecular and Solid State Physics


### Atomic form factors

In the range of scattering vectors between  $0 < q < 25 \text{ \AA}^{-1}$ , the atomic form factor is well approximated by the expression, [1]

$$f(q) = \sum_{i=1}^4 a_i \exp\left(-b_i \left(\frac{q}{4\pi}\right)^2\right) + c,$$

where the values of  $a_i$ ,  $b_i$ , and  $c$  are tabulated below. The different atomic form factors for the elements can be plotted using the form below.

Atomic form factor for H



Element	$a_1$	$b_1$	$a_2$	$b_2$	$a_3$	$b_3$	$a_4$	$b_4$	$c$
H	0.489918	20.6593	0.262003	7.74039	0.196767	49.5519	0.049879	2.20159	0.001305
H1-	0.897661	53.1368	0.565616	15.187	0.415815	186.576	0.116973	3.56709	0.002389
He	0.8734	9.1037	0.6309	3.3568	0.3112	22.9276	0.178	0.9821	0.0064



### Primitive lattice vectors:

$$\begin{aligned}\vec{a}_1 &= 4.12\text{E-10} \hat{x} + 0 \hat{y} + 0 \hat{z} \text{ [m]} \\ \vec{a}_2 &= 0 \hat{x} + 4.12\text{E-10} \hat{y} + 0 \hat{z} \text{ [m]} \\ \vec{a}_3 &= 0 \hat{x} + 0 \hat{y} + 4.12\text{E-10} \hat{z} \text{ [m]}\end{aligned}$$

### Basis:

The positions of the atoms are given in fractional coordinates between -1 and 1.

Cs1+	0	$\vec{a}_1 +$	0	$\vec{a}_2 +$	0	$\vec{a}_3$
Cl1-	0.5	$\vec{a}_1 +$	0.5	$\vec{a}_2 +$	0.5	$\vec{a}_3$
		$\vec{a}_1 +$		$\vec{a}_2 +$		$\vec{a}_3$
		$\vec{a}_1 +$		$\vec{a}_2 +$		$\vec{a}_3$
		$\vec{a}_1 +$		$\vec{a}_2 +$		$\vec{a}_3$

submit

Al (fcc)

NaCl (fcc)

CsCl (sc)

SrTiO3 (sc)

GaAs (Zincblend, fcc)

GaN (Wurtzite, hex)

### Primitive reciprocal lattice vectors

$$\begin{aligned}\vec{b}_1 &= 2\pi \frac{\vec{a}_2 \times \vec{a}_3}{\vec{a}_1 \cdot (\vec{a}_2 \times \vec{a}_3)} = 1.525\text{e+10} \hat{k}_x + 0.000 \hat{k}_y + 0.000 \hat{k}_z \text{ [m}^{-1}\text{]} \\ \vec{b}_2 &= 2\pi \frac{\vec{a}_3 \times \vec{a}_1}{\vec{a}_1 \cdot (\vec{a}_2 \times \vec{a}_3)} = 0.000 \hat{k}_x + 1.525\text{e+10} \hat{k}_y + 0.000 \hat{k}_z \text{ [m}^{-1}\text{]} \\ \vec{b}_3 &= 2\pi \frac{\vec{a}_1 \times \vec{a}_2}{\vec{a}_1 \cdot (\vec{a}_2 \times \vec{a}_3)} = 0.000 \hat{k}_x + 0.000 \hat{k}_y + 1.525\text{e+10} \hat{k}_z \text{ [m}^{-1}\text{]}\end{aligned}$$

### Structure factors

The value of  $|n_{\vec{G}}|$  for the 000 diffraction peak is the total number of electrons in the primitive unit cell. The intensities of the peaks in an x-ray diffraction experiment

$hkl$	$ \vec{G}  \text{ \AA}^{-1}$	$ n_{\vec{G}} $	$ n_{\vec{G}} ^2$	$\text{Re}\{n_{\vec{G}}\}$	$\text{Im}\{n_{\vec{G}}\}$
000	0.000	72.00	5184	72.00	0.000
-100	1.525	34.43	1185	34.43	5.333e-8
0-10	1.525	34.43	1185	34.43	5.333e-8
00-1	1.525	34.43	1185	34.43	5.333e-8
001	1.525	34.43	1185	34.43	-5.333e-8



# Structure factor

---

$$n_{\vec{G}} = \frac{1}{V} \sum_j f_j(G) e^{-i\vec{G} \cdot \vec{r}_j}$$

A structure factor is the Fourier transform of the electron density of the basis of a crystal evaluated at a reciprocal lattice vector. Since the electron density of the basis can be approximated as a sum over the electron densities of the atoms in the basis, the Fourier transform of the electron density of the basis is a sum of the Fourier transforms of the electron densities of the atoms in the basis.



An x-ray experiment measures the scattered intensity  $|F_G|^2$ . The phase information is lost. This is proportional to  $|n_G|^2$ .

# crystal structure solution

---


## **structural solution of the DNA**

F.Crick, J.Watson, M.Wilkins  
nobel laureate 1962 for medicine



Rosalind Franklin

# crystal structure solution



"Guess" the crystal structure





Table 4.3.2.2 | pdf |

Elastic atomic scattering factors of electrons for neutral atoms and s up to  $2.0 \text{ \AA}^{-1}$

Element	Z	$a_1$	$a_2$	$a_3$	$a_4$	$a_5$	$b_1$	$b_2$	$b_3$
H	1	0.0349	0.1201	0.1970	0.0573	0.1195	0.5347	3.5867	12.3471
He	2	0.0317	0.0838	0.1526	0.1334	0.0164	0.2507	1.4751	4.4938
Li	3	0.0750	0.2249	0.5548	1.4954	0.9354	0.3864	2.9383	15.3829
Be	4	0.0780	0.2210	0.6740	1.3867	0.6925	0.3131	2.2381	10.1517
B	5	0.0909	0.2551	0.7738	1.2136	0.4606	0.2995	2.1155	8.3816
C	6	0.0893	0.2563	0.7570	1.0487	0.3575	0.2465	1.7100	6.4094
N	7	0.1022	0.3219	0.7982	0.8197	0.1715	0.2451	1.7481	6.1925
O	8	0.0974	0.2921	0.6910	0.6990	0.2039	0.2067	1.3815	4.6943
F	9	0.1083	0.3175	0.6487	0.5846	0.1421	0.2057	1.3439	4.2788
Ne	10	0.1269	0.3535	0.5582	0.4674	0.1460	0.2200	1.3779	4.0203
Na	11	0.2142	0.6853	0.7692	1.6589	1.4482	0.3334	2.3446	10.0830
Mg	12	0.2314	0.6866	0.9677	2.1882	1.1339	0.3278	2.2720	10.9241
Al	13	0.2390	0.6573	1.2011	2.5586	1.2312	0.3138	2.1063	10.4163
Si	14	0.2519	0.6372	1.3795	2.5082	1.0500	0.3075	2.0174	9.6746
P	15	0.2548	0.6106	1.4541	2.3204	0.8477	0.2908	1.8740	8.5176
S	16	0.2497	0.5628	1.3899	2.1865	0.7715	0.2681	1.6711	7.0267
Cl	17	0.2443	0.5397	1.3919	2.0197	0.6621	0.2468	1.5242	6.1537
Ar	18	0.2385	0.5017	1.3428	1.8899	0.6079	0.2289	1.3694	5.2561

Compare  $|S_G|^2$  to the measurements



From the atomic form factors, calculate the structure factors  $n_G$ .

$$S_{\vec{G}} = \sum_j f_j(G) e^{-i\vec{G} \cdot \vec{r}_j}$$

Sum over basis

position of atom  $j$  of the basis